GEOLOGIC SETTING The Tycho quadrangle is centrally located in the southern half of the Earthside hemisphere of the Moon. The area is characterized by a high density of craters, the largest of which-Stöfler-is about 140 km in diameter. The northern and eastern parts of the quadrangle are dominated by plains (pIp, Ip)

and hilly terra (IpIt) units of regional extent, and the western part by the crater Tycho and its ejecta blanket. Structural fea-

tures probably associated with the Imbrium basin to the north

and the Orientale and Humorum basins to the northwest are

present, although blanketing units related to these basins are

STRATIGRAPHY

Pre-Imbrian plains-forming material (pIp).—Parts of the

quadrangle are occupied by densely cratered, relatively bright

blanketing plains material of pre-Imbrian age. The unit shares

some of the characteristics of both hilly terra (IpIt) and Imbrian plains (Ip) units; it is apparently intergradational with them in some areas and is separable from them only with difficulty. As mapped, pre-Imbrian plains occupy broad shallow depressions in the north-central, eastern, and south-central parts of the quadrangle. The unit lacks the hummocky topography

and lineated texture of the hilly terra materials, but has the

same albedo. Also, like the Imbrian plains unit, it is apparently

of volcanic origin and may have accumulated as a series of

highly fluid ash flows—probably from several centers of erup-

Pre-Imbrian and Imbrian (?) hilly terra material (IpIt).

Much of the terra in the quadrangle that is not occupied by

mapped craters is relatively flat but has a gently hilly, rolling,

or finely to irregularly hummocky or faintly lineated surface.

Much of this surface texture is believed to be intrinsic to a

material stratigraphic unit, mapped as hilly terra material, that

fills the intercrater areas. In the east half of the quadrangle

this unit covers most of the rim deposits of lower pre-Imbrian

craters (unit pIcr₁) as well as broad expanses having low crater

densities. In areas where the finely hummocky topography

which best characterizes this unit is absent, rim deposits of

middle and upper pre-Imbrian craters (units pIcr₂ and pIcr₃)

are well preserved and have distinct positive relief. Hilly ma-

terial is interpreted to be volcanic in origin and may be com-

posed of ash-flow deposits of variable thickness. Inasmuch as topographically distinguishable sources are not present in the

area, the volcanic materials may be derived from some of the

small craters that are so numerous here (see also Offield, 1971).

Part of the unit is probably composed of lunar regolith devel-

not recognized.

I-713 (LAC 112)



Slope material Material on steep slopes, primarily crater walls, distinctly brighter than Tycho ray material; albedo 0.150-0.220 (throughout explanation, albedo data from Pohn and Wildey, 1970) Bedrock and blocky talus exposed by impact, faulting, and mass wasting. Generally younger than Tycho ray material

Ccfs, floor material, smooth. Few hummocks. Trav-

ersed by numerous fissures tens of meters wide

barely visible at Orbiter IV resolution (about 100

m) but apparent on Orbiter V photographs (reso-

lution 4 m). Density of craters < 1 km diameter

Cefh, floor material, hummocky. Broadly rounded

sures tens of meters wide. Density of craters < 1

Ccrb, rim material, blocky. Coarse and rather angu-

Ccrh, rim material, hummocky. Large subdued hum-

mocks, about 5 km diameter, and subdued ridges

subconcentric to rim crest. Heavily lineated in

part. Distal to unit Cerb where that unit present

Cerr, rim material, radial. Continuous surface hav-

ing ropy or braided texture radial to Tycho. Sub-

dued satellitic craters abundant (but not mapped)

near outer contact; grades outward into facies dom-

Ccw, wall material. Finely hummocky material on

terraced wall. Small irregular and locally sinu-

ous ridges and rimmed valleys trend down steep

Ccp, peak material. Sharply angular complex of

inated by satellitic craters (unit Csc)

ridges and peaks near center of crater

lar hummocks about ½ km diameter adjacent to

hummocks 1/4 to 3 km diameter. Numerous fis-

lower than on crater rim

km lower than on rim



Ccfs, post-crater lava or other volcanic filling sub-

sequently cut by tension cracks as result of cooling

may be volcanic domes. Fissures possibly caused

ty coalescence of individual volcanic elements. Some

Ccfh. post-crater viscous volcanic filling. Hummocks

Ccrb, impact ejecta consisting mainly of large jumbled

Ccrh, impact ejecta covered by moderately thin blan-

ket of fine material deposited from base surge

charnels and ridges possibly formed by downslope

movement of particulate material while crater in-

Ccp, part of lens of intensely brecciated material up-

lifted immediately after excavation of crater; pos-

terior still hot from impact shock melting

craters in floor may be effusion vents

blocks swept clean of base-surge deposits

Ccrr, impact ejecta deposited by base surge Ccw, slumped rim materials and talus. Radial

sibly later volcanic material in part

ameter) for subdivision of units

lution. Forms flat floor in Kaiser C

and microneteorite bombardment

Interpretation

Materials of crater Tycho

EXPLANATION

Cc, crater material, undivided. Material of two bright, sharprimmed craters in west-central part of quadrangle too small (<3~km~rim~crest~diameter) to distinguish subunits. Superposed on Tycho materials $Ccr, rim\ material.$ Surrounds single crater (Hell QA) in north-central part of quadrangle; crater has sharp rim and very bright rays which interrupt Tycho rays Interpretation Moterials of impact craters younger than Tycho



Streaks and halos having higher albedo than Tycho rays. Occurs around three small (<5~km) Copernican craters and in east-west rending patch on floor of Walter Fine-grained ejecta of small Copernican impact craters; in Walter, bright slope material and ejecta of secondary impact craters of unknown provenance NOTE: Ray material from Tycho covers most of quadrangle



and has been omitted to clarify map and to show rays of

Characteristics Material of small generally sharp-rimmed craters in clusters and strings subradial to crater Tycho; craters are elongate or round single craters or coalescing craters. Many strings exhibit a herringbone or braided pattern splaying outward

Tycho but recur on floors and opposite walls, indicating ballistic shalowing—ejecta impacted surface at angle less than slope of walls




from Tycko Material of secondary impact craters produced by ejecta from Tycho. Some patches stop abruptly at crater rim crests facing

oped in place.

Imbrian plains-forming material (Ip).—Mare materials are absent in the quadrangle; however, a plains-forming unit of Imbrian age, similar in some respects to mare material, is present. It occupies the floors of many pre-Imbrian and early Imbrian craters as well as other topographic lows in all parts of the quadrangle, except in the immediate vicinity of the crater Tycho. The unit has about the same density of superposed craters as some mare units mapped elsewhere on the lunar surface but has a slightly higher albedo than typical mare material. On the other hand, it is less densely cratered and slightly darker than the pre-Imbrian plains-forming unit. Its presence in craters as young as earliest Imbrian and absence in younger craters suggest that it probably was emplaced in early Imbrian time. Crater materials.—The relative ages of crater materials are determined by a comparison of their morphology within selected size categories (Pohn and Offield, 1970), and their assignment to the lunar time-stratigraphic systems follows Offield (1971). Rim and floor materials of the crater Tycho. - Tycho is the youngest large crater on the near side of the Moon and is generally recognized to have formed by impact. Because of its youthfulness, the features characteristic of impact-formed craters are well developed. Of particular interest are the materials that apparently accumulated by flowage and occupy much of the floor and smaller areas of the crater rim. Their mode of origin is controversial. In describing the geology of the Surveyor VII landing site, Shoemaker and others (1968, p. 26), for

example, give three plausible explanations for the origin of flows on the rim of Tycho: (1) The flows are volcanic and have been produced by extrusion of lava derived from depth, possibly formed or extruded in response to the forces that produced the crater. (2) The flows are relatively cold debris flows, which were mobile or fluid because they contained either liquid water or gas. (3) The flows are relatively hot debris flows, which contained molten, or partly molten, rock (and possibly some gas) heated by shock and by viscous flow during ejection of the debris from



STRUCTURE AND MATERIALS OF CRATER CHAINS AND CLUSTERS An analysis of lineaments from Lunar Orbiter IV photographs shows strong directional preference for N. 35° E., N. 40° W., and N. 60° W. trends. On this part of the lunar surface none of these directions is radial to the major lunar basins. On the other hand, an analysis of the major fault directions shows pronounced radiality to the nearby basins, and to the more distant Imbrium basin (fig. 1). In addition, many of the long axes of the crater chains and some clusters (whose materials are mapped as units Ich, Icc, and Iccr) are subradial to the Imbrium basin. These subradial chains could be volcanic in origin, but it seems more likely that they are secondary impact craters of

the Imbrium basin. Many of the chain craters have pronounced rim deposits and shapes which resemble those of secondary impact craters mapped elsewhere on the Moon. Their detailed morphology is similar to that of the oldest craters on the earliest Imbrian deposits of the Apennine Mountains. Thus, they probably are secondary craters of early Imbrian age. GEOLOGIC HISTORY The earliest decipherable events that occurred in the quadrangle were the formation in pre-Imbrian time (presumably by impact) of many of the large craters. Some volcanism also

occurred in the late pre-Imbrian, resulting in the deposition of plains and, possibly, hilly terra units. The impact which formed the Imbrium basin (centered about 1,200 miles north of the Tycho quadrangle) in earliest Imbrian time ejected enormous volumes of material, some of which apparently formed secondary impact crater chains and clusters within the quadrangle. Widespread volcanism occurred during the Imbrian Period, and in early Imbrian time, another plains-forming unit was deposited. A few impact craters were formed during the Eratosthenian Period. Some impact craters were probably formed early in the Copernican Period, but the evidence for dating these craters (ray material) has been largely obscured by the rays from Tycho. Formation of the crater Tycho in late Copernican time is the most recent major event in the area. Deposits formed by the Tycho impact (unit Ccrr) subdued the topography of the southwestern part of the quadrangle, and abundant secondary craters (Csc) were formed in the remainder of the area. In addition, the quadrangle and much of the near side of the Moon was mantled with thin discontinuous Tycho ray deposits (not mapped).

REFERENCES

Offield, T. W., 1971, Geologic map of the Schiller quadrangle of the Moon: U.S. Geol. Survey Misc. Geol. Inv. Map I-691. Pohn, H. A., and Offield, T. W., 1970, Lunar crater morphology and relative-age determination of lunar geologic units-Pt. 1 Classification, in Geological Survey research 1970: U.S. Geol. Survey Prof. Paper 700-C, p. C153-C162. Pohn, H. A., and Wildey, R. L., 1970, A photoelectric-photographic study of the normal albedo of the Moon: U.S. Geol. Survey Prof. Paper 599-E, p. E1-E20. Shoemaker, E. M., Batson, R. M., Holt, H. E., Morris, E. C., Rennilson, J. J., and Whitaker, E. A., 1968, Television observations from Surveyor VII, in pt. II, Science results, of Surveyor VII mission report: California Inst. Technology, Jet Propulsion Lab. Tech. Rept. 32-1264, p. 23-44.

Cratered dome material Characteristics Domes with small summit pits. Circular in plan, range from 2 to 10 km diameter, convex upward. Surfaces generally smooth but finely braided texture in places Interpretation

Probably volcanic domes

LUNAR ORBITER PHOTOGRAPHIC COVERAGE OF TYCHO QUADRANGLE

LO V H 125-128, Orbiter V high-resolution frames (approx. 8 m

identification resolution); LO V M125-128, Orbiter V medium reso-

lution frames (approx. 60 m); all other numbers refer to Orbiter IV

high-resolution frames (approx. 100 m)

Chain crater materials

Characteristics

approximately located; short-dashed where gradationalConcealed contact

Inferred fault or fault scarp



GEOLOGIC MAP OF THE TYCHO QUADRANGLE OF THE MOON

1972

features portrayed.

INDEX MAP OF THE EARTHSIDE HEMISPHERE OF THE MOON

Number above quadrangle name refers to lunar base chart (LAC series); number below refers to published geologic map

Materials of crater chains and clusters

lescent elongate craters about 2 to 10 km diameter, generally decreasing in diameter along length of chain; normally no divid-Icc, crater cluster material. Similar to units too small or poorly defined for mapchains except craters arranged randomly Iccr, cluster rim materials. Moderately hummocky rims, concave upward; more conspicuous on one side of cluster than the other

(Icr), wall (Icw), and floor (Icf) materials Interpretation Materials of impact cravers of indeterminate Imbrian age

craters but more subdued. Morphology and (or) superposition relations indicate age

older than Eratosthenian but younger than

late pre-Imbrian. In craters < 7 km sub-

ping; in larger craters divided into rim

Interpretation Probably most are secondary impact craters of Imbrium basin (see text)

more subdued; in addition to abundant

small craters, has at least one large crater Icw1, wall material. Walls terraced but considerably subdued and coalesced Icf₁, floor material. Similar to unit Icf₂ Icp₁, peak material. Similar to unit Icp₂ Interpretation Same as for upper Imbrian crater materials but craters more subdued and therefore older

Crater materials

strongly subdued and crenulated

Icr₁, rim material. Similar to unit Icr₂ but

Ec, crater materials, undivided. Fresh nonrayed craters too small (<5 km rimcrest di-

Ecr, rim material. Positive relief outside rim crest; concave upward; hummocky near

Ecw, wall material. Steep walls in craters; appears finely granular at Orbiter IV reso-

Materials of impact craters more degraded than Copernican materials by seismic shaking

rim crest grading outward into subradial and braided texture

Hilly and furrowed material Characteristics

Ic2, crater materials, undivided. Rim crests moderately subdued

Icw2, wall material. Walls terraced but not coalesced

Icp₂, peak material. Similar to unit Ccp but more subdued

 ${
m Icr}_2$, rim material. In craters > 20 km, similar to ${
m Ecr}$ but has abundant craters < 3 km

Icf₂, floor material, undivided. Mixture of smooth and hummocky floor materials

Icfs, floor material, smooth. Similar to unit Ccfs but without fissures or hummocks

Materials of impact craters similar in origin to those of younger craters but more highly

Icfh, floor material, hummocky. Similar to unit Ccfh but more subdued

modified by impact, volcanism, mass-wasting, and seismic shaking

Material of low, circular to elongate gently Ic₁, crater materials, undivided. Rim crests rounded hills and subdued northeast-trending ridges; in places small unmapped crater chains occur between ridges. Gradational with hilly terra material. Blankets rim B but other Imbrian craters superposed Volcanic material of Imbrian age

Smooth plains material

sode of volcanism

Interpretation Volcanic materials, possibly ash-flow tuffs

Plains material

Mainly in low intercrater areas. Flat to

gently undulatory; generally more densely

cratered than hilly terra material and

younger plains material but locally inter-

gradational with these units. Partly covers

Blanket of material thick enough to obscure

canic materials, most likely ash flow tuffs;

pre-existing topography; probably fluid vol-

in part may be deep regolith developed in

some pre-Imbrian crater rim materials

Interpretation

Flat cratered surfaces in many crater floors and other depressions. Approximately level within a depression; elevation varies from depression to depression. Crater density approximately same as or somewhat higher than on mare materials (not in quadrangle) Queried near crater Tycho where numerous satellitic craters obscure characteristics and interrupt lateral continuity of unit

emplaced from many separate vents during

major, probably extended, early Imbrian epi-

Crater materials

pIc, crater materials undivided. Subunits

not mapped because of burial or small

crater size. Rims markedly to completely

pIcr, rim material. Rim crests subdued and

finely crenulated. Rims commonly inter-

plew, wall material. Similar to units plew,

Same as for other pre-Imbrian craters but

classification within pre-Imbrian uncertain

rupted by large craters

plew2, and plew3

Interpretation

burial or small crater size. Rims markedly subdued and rounded, partly sharp craters (< 20 km); poorly developed in larger craters finely granular at Orbiter IV resolution

Crater materials pIc2, crater materials undivided. Similar to unit pIc3 but markedly to completely subdued plcr₂, rim material. Subdued and highly cratered; rim crest mostly crenulated; no sharp parts pIcw2, wall material. Hummocky; terraces subdued and barely discernible; radial channels well developed pIcf₂, floor material. Similar to but more subdued than unit pIcf₃ pIcp₂, peak material. Densely cratered highly subdued peaks in Walter; buried in Huggins. No sharp crests; slopes gentle

Characteristics pIcr rim material. Highly subdued to barely discernible rims and rim crests; rim crest discontinuous; heavily cratered pIcw₁, wall material. Hummocky; slopes shallow; few terrace segments Interpretation Same as for middle and upper pre-Imbrian crater materials.


ch, crater material. Chains <20 km long of coalescent, elongate craters. Crater diameters decrease toward one end of chain. No apparent dividing septa chr, rim material. Low slightly hummocky rims surrounding the chains Interpretation Chains of volcanic craters, probably exploContact

Buried unit indicated by symbol in parentheses

sion vents: rim material composed of pyro- Long dashed where approximately located; dotted where concealed. Line at base of scarp; bar and ball on apparent downthrown side. Fault either offsets unit or forms scarp against which younger rocks were de-

Narrow trough _____ Lineament Gentle linear depression.

Interpretation: Fault, buried or exposed sense of movement undetermined __.._.._.._ Buried rim crest Depression



mocky and pitted at 1-2 km scale; locally rough; rough and highly lineated between craters Nonius and Fernelius. Ob scures early pre-Imbrian crater rim materials and present on wall and floor materials; also present on some middle and upper pre-Imbrian rim materials. Queried where patches small or blurred by Tycho ejecta Interpretation Probably thin to moderately thick blanket, possibly of ash-flow and ash-fall tuff, deposited on pre-existing terra and crater

materials. Lineaments structural or depositional or both. Apparent absence on middle and upper pre-Imbrian crater walls possibly due to downslope movement. Age probably pre-Imbrian but possibly Imbrian in places

pIc3, crater materials undivided. Subunits not mapped because of pIcr₂, rim material. Subdued raised rims; rim crest partly crenulated plew3, wall material. Moderately steep to subdued, hummocky terraced walls having some radial channels. Channels well developed in small plcf₃, floor material. Low hills, ridges and broad hummocks; appears plcp3, peak material. Moderately cratered, few sharp ridge crests,

slopes gentle to steep. Moderately subdued peaks in Licetus Interpretation Origin similar to corresponding younger crater units but more highly modified by impact, volcanism, mass-wasting, and seismic-shaking

Interpretation Same as for upper pre-Imbrian crater materials

Crater materials

clastic ejecto and possibly some country rock. Age not determined

For sale by U.S. Geological Survey, price \$1.00